arXiv:math-ph/0201028v2 21 Jan 2002

NORM ESTIMATES OF ALMOST MATHIEU OPERATORS

FLORIN P. BOCA AND ALEXANDRU ZAHARESCU

Abstract. We estimate the norm of the almost Mathieu operator $H_{\theta,\lambda} = U_\theta + U_\theta^* + \frac{\lambda}{2}(V_\theta + V_\theta^*)$, viewed as an element in the rotation C^*-algebra $A_\theta = C^*(U_\theta, V_\theta \text{ unitaries}\,;\, U_\theta V_\theta = e^{2\pi i\theta} V_\theta U_\theta)$. In the process, we prove the inequality

$$\|H_{\theta,\lambda}\| \leq \sqrt{4 + \lambda^2 - \left(1 - \frac{1}{\tan \pi\theta}\right)\left(1 - \sqrt{\frac{1 + \cos^2 4\pi\theta}{2}}\,\right)\min(4, \lambda^2)}$$

for every $\lambda \in \mathbb{R}$ and every $\theta \in [1/4, 1/2]$. This significantly improves the inequality $\|H_{\theta,2}\| \leq 2\sqrt{2}$, $\theta \in [1/4, 1/2]$, conjectured by Béguin, Valette and Zuk.

1. Introduction and Statement of Results

An almost Mathieu operator is a discrete Schrödinger operator that acts on the Hilbert space $\ell^2 = \ell^2(\mathbb{Z})$ as

$$H(\theta, \lambda, \phi)\xi_n = \xi_{n+1} + \xi_{n-1} + \lambda \cos 2\pi(n\theta + \phi)\, \xi_n,$$

where θ, λ and ϕ are real numbers, and $(\xi_n)_n$ denotes the canonical orthonormal basis in ℓ^2. The study of the spectral properties of this class of operators has attracted a significant amount of interest in the past couple of decades (see [5], [3], [8], [1], [10], [14], [12], [7] for some of the most important developments). Most of this work has focused on the "Ten Martini problem" of M. Kac, concerning the possible values of the labels of the gaps that appear in the spectrum of these kinds of operators, and on the localization properties of the spectrum.

The almost Mathieu operator $H(\theta, \lambda, \phi)$ can be regarded as the image of the self-adjoint element $H_{\theta,\lambda} = U + U^* + (\lambda/2)(V + V^*)$ in the representation of the rotation C^*-algebra $A_\theta = C^*(U, V \text{ unitaries}\ ;\ UV = e^{2\pi i\theta} VU)$ that maps U to the bilateral shift u_0 defined on ℓ^2 by $u_0\xi_n = \xi_{n-1}$, and V to the diagonal unitary v_0 defined by $v_0\xi_n = e^{2\pi i(n\theta+\phi)}\xi_n$. The operator $H_\theta = H_{\theta,2}$ is called a Harper operator.

When $\theta = p/q$ is rational with $0 \leq p < q$ coprime integers, the spectrum of $H_{\theta,\lambda}$, viewed as an element of A_θ, consists either in the union of q disjoint intervals (when q is odd), or of $q - 1$ disjoint intervals (when q is even). This is best illustrated in Hofstadter's butterfly ([11]) in Figure 1.

Received by the editors February 7, 2008.



FIGURE 1. The spectra of the Harper operators $H_{p/q}$ for $0 \leq p/q \leq 1$

The irreducible representations of $A_{p/q}$ can be classified up to unitary equivalence. They all have dimension q and are given by

$$\pi_{z_1,z_2}(U) = z_1 U_0, \qquad \pi_{z_1,z_2}(V) = z_2 V_0$$

for $z_1, z_2 \in \mathbb{T}$, where

$$U_0 = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & 0 & \dots & 0 & 0 \end{pmatrix}, \quad V_0 = \begin{pmatrix} 1 & & & & \\ & e^{2\pi i\theta} & & & \\ & & e^{4\pi i\theta} & & \\ & & & \ddots & \\ & & & & e^{2\pi i(q-1)p/q} \end{pmatrix}$$

and $\theta = \dfrac{p}{q}$.

We also have in this case (θ rational)

$$\|H_{\theta,\lambda}\| = \|h_{\theta,\lambda}\|, \tag{1.1}$$

where

$$h_{\theta,\lambda} = U_0 + U_0^* + \frac{\lambda}{2}(V_0 + V_0^*) \in M_q(\mathbb{C}) = \mathcal{B}(\ell^2(\mathbb{Z}_q)). \tag{1.2}$$

The map $\theta \mapsto \|H_{\theta,\lambda}\|$ is $\mathrm{Lip}^{1/2}$ continuous on $[0,1]$ (see [1], [4], [9] or [13] for different proofs with various degrees of generality).

All these properties of rotation C^*-algebras and of almost Mathieu operators can be found, with self-contained proofs, in the first four chapters of [6].

The first estimates on the norm of almost Mathieu operators were given by C. Béguin, A. Valette and A. Zuk ([2]), who proved the inequality

$$\|H_\theta\| \leq 2(1+\sqrt{2}+\cos 2\pi\theta) \qquad \theta \in [0,1]. \tag{1.3}$$

This is effective only in a neighbourhood of $\theta = 1/2$, as the norm of H_θ is no greater than 4.

It was also conjectured by these authors in [2] that[1]

$$\|H_\theta\| \leq 2\sqrt{2} \qquad \theta \in [1/4, 1/2]. \tag{1.4}$$

A proof of (1.4) has appeared in [15]. Unfortunately, the trigonometric formulas at page 157 are not correct and relation (13) in [15] should change to

$$(\sqrt{2}+s_n-\gamma_n)(\sqrt{2}-s_n+\gamma_{n-1}) \geq 1.$$

This makes the forthcoming arguments at pages 158-159, concerning the possibility of chosing γ_n such that this inequality be satisfied, to be incorrect.

The original motivation of this work was to give a complete proof of the inequality (1.4). This task is achieved first in Section 2, where we prove for every $\lambda \in \mathbb{R}$ and every $\theta \in [1/4, 1/2]$ the inequality

$$\|H_{\theta,\lambda}\| \leq \sqrt{4+\lambda^2}\,. \tag{1.5}$$

Note that the equality holds in (1.5) at $\theta = 1/4$ and $\theta = 1/2$ for any λ. In this section we also prove in the range $\theta \in [0, 1/2]$ the estimate

$$\|H_{\theta,\lambda}\| \leq M_\lambda(\theta) = \sqrt{4+\lambda^2+4|\lambda|(\cos\pi\theta-\sin\pi\theta)\cos\pi\theta}. \tag{1.6}$$

In Section 3, we further sharpen the upper bound $\sqrt{4+\lambda^2}$ for $\|H_{\theta,\lambda}\|$ on $[1/4, 1/2]$, and prove for every $\theta \in [1/4, 1/2]$ the inequality

$$\begin{aligned} \|H_{\theta,\lambda}\| &\leq M_\lambda(\theta) \\ &= \sqrt{4+\lambda^2-\left(1-\frac{1}{\tan\pi\theta}\right)\left(1-\sqrt{\frac{1+\cos^2 4\pi\theta}{2}}\right)\min(4,\lambda^2)}, \end{aligned} \tag{1.7}$$

which provides, when $\lambda = 2$, a significant improvement of (1.4) (see Figures 3 and 2). In the case $\lambda = 2$, the upper bound estimates (1.6) and (1.7) are compared in Figure 2 with (1.3) and with the bound

$$\|H_\theta\| \leq \begin{cases} 2+2\cos\pi\theta & \text{if } \sin^2\pi\theta \leq (\sqrt{5}-1)/2 \\ 2\sqrt{1+1/\sin^2\pi\theta} & \text{if } \sin^2\pi\theta \geq (\sqrt{5}-1)/2, \end{cases} \tag{1.8}$$

proved (correctly) in [15].

In the last part of the paper, we give some explicit lower bounds for the norm of Harper operators, proving the inequality

$$\|H_\theta\| \geq m(\theta) = \max\big(f_1(\theta), f_2(\theta), f_3(\theta)\big) \qquad \theta \in [0, 1/2], \tag{1.9}$$

---

[1] The existence of the $*$-antiautomorphism of A_θ that maps U to U^* and V to V^* implies that the spectra of $H_{\theta,\lambda}$ and $H_{1-\theta,\lambda}$ coincide for any $\theta \in [0,1]$ and any $\lambda \in \mathbb{R}$.

where $f_1(\theta), f_2(\theta), f_3(\theta) \geq 0$ are given by

$$\begin{aligned} f_1(\theta)^2 = 6 - {} & \frac{4}{1 + \frac{1}{\sin \pi\theta} + \sqrt{1 + \frac{1}{\sin \pi\theta}}} \\ & + \frac{2}{1 + 4 \sin \pi\theta \cos^2 \pi\theta} + \frac{8 \cos^2 \pi\theta}{(1 + \sin \pi\theta)^{3/2}}, \end{aligned} \tag{1.10}$$

$$\begin{aligned} & f_2(\theta)^2 = 4 + \frac{2}{\sqrt{1 + |\sin 4\pi\theta|}} + \frac{2 \cos^2 2\pi\theta}{1 + |\sin 4\pi\theta|} \\ & + 2\sqrt{\left(\frac{\sin^2 2\pi\theta}{1 + |\sin 4\pi\theta|} \right)^2 + \frac{16 \cos^4 \pi\theta}{(2 + |\tan 2\pi\theta|)^2} \Big(1 + \frac{\cos 2\pi\theta}{\sqrt{1 + |\sin 4\pi\theta|}} \Big)^2}, \end{aligned} \tag{1.11}$$

$$\begin{aligned} & f_3(\theta)^2 = 4 + \frac{4}{5} (\cos 2\pi\theta + 2\cos^2 2\pi\theta + 2 \cos^4 2\pi\theta) \\ & + \sqrt{\left(2 - \frac{4(\cos 2\pi\theta + 2\cos^2 2\pi\theta + 2\cos^4 2\pi\theta)}{5} \right)^2 + \left(\sqrt{10} + \frac{8 \cos 2\pi\theta}{\sqrt{10}} \right)^2}. \end{aligned} \tag{1.12}$$

In particular we get

$$\min_{1/4 \leq \theta \leq 1/2} \|H_\theta\| \geq \min_{1/4 < \theta < 1/2} f_1(\theta) \approx \sqrt{6.59303} \approx 2.56769. \tag{1.13}$$

This is a good estimate for $\min\limits_{1/4 \leq \theta \leq 1/2} \|H_\theta\|$, which appears to be, by numerical computations, just fractionally larger than 2.59.

We also get (see Figure 3)

$$\min_{0 \leq \theta \leq 1/4} \|H_\theta\|^2 \geq 7.82387 \tag{1.14}$$

and for all $0 \leq \theta \leq 0.23441$ we get

$$\|H_\theta\|^2 \geq 8. \tag{1.15}$$

Our estimates (1.6), (1.7) and (1.9) on the norm of the Harper operator H_θ are illustrated in Figure 3.

Numerical computations appear to indicate that inequality (1.15) might hold for every $\theta \in [0, 1/4]$. It would be interesting to clarify this point.

2. A PROOF OF THE INEQUALITY $\|H_{\theta,\lambda}\| \leq \sqrt{4 + \lambda^2}$, $1/4 \leq \theta \leq 1/2$

Throughout this section, we take $\theta = p/q$, denote

$$C_n = C_n(\theta) = \cos 2n\pi\theta \qquad n \in \mathbb{Z},$$

and consider the self-adjoint $q \times q$ matrix $h_{\theta,\lambda}$ defined in (1.2). That is, $h_{\theta,\lambda}$ acts on $\ell^2(\mathbb{Z}_q)$ by

$$h_{\theta,\lambda} \epsilon_n = \epsilon_{n+1} + \epsilon_{n-1} + \lambda C_n \epsilon_n \qquad n \in \mathbb{Z}_q,$$

where $(\epsilon_n)_{n \in \mathbb{Z}_q}$ denotes the canonical orthonormal basis in $\ell^2(\mathbb{Z}_q)$.

In this section and the next one, we set $\sum\limits_m = \sum\limits_{m \in \mathbb{Z}_q}$.



FIGURE 2. The upper bound estimates (1.3),(1.8),(1.6) and (1.7)



FIGURE 3. The graphs of the functions $[0, 1/2] \ni \theta \mapsto M_2(\theta), \|H_\theta\|, m(\theta)$

Let $(X_m)_{m\in\mathbb{Z}_q}$ be a unit eigenvector in $\ell^2(\mathbb{Z}_q)$ for the eigenvalue E of $h_{\theta,\lambda}$ with $X_m = X_m(\theta) \in \mathbb{R}$. Then

$$X_{m+1} + X_{m-1} + \lambda C_m X_m = E X_m \qquad m \in \mathbb{Z}_q, \tag{2.1}$$

which gives

$$\begin{aligned}
E^2 &= \sum_m (EX_m)^2 = \sum_m (X_{m+1} + X_{m-1} + \lambda C_m X_m)^2 \\
&= 2 + \lambda^2 \sum_m C_m^2 X_m^2 + 2 \sum_m X_{m+1} X_{m-1} \\
&\quad + 2\lambda \sum_m X_m X_{m-1} (C_m + C_{m-1}) \\
&= 2 + \lambda^2 \sum_m (1 - \sin^2 2m\pi\theta) X_m^2 + 2 - \sum_m (X_{m+1} - X_{m-1})^2 \\
&\quad + 2\lambda \sum_m X_m X_{m-1} (C_m + C_{m-1}) \\
&= 4 + \lambda^2 - \sum_m (X_{m+1} - X_{m-1})^2 - \lambda^2 \sum_m X_m^2 \sin^2 2m\pi\theta \\
&\quad + 2\lambda \sum_m X_m X_{m-1} (C_m + C_{m-1}).
\end{aligned} \tag{2.2}$$

It also follows from (2.1) that

$$\begin{aligned}
\lambda \sum_m X_m X_{m-1} C_{m-1} &= \sum_m X_m (EX_{m-1} - X_m - X_{m-2}) \\
&= E \sum_m X_m X_{m-1} - 1 - \sum_m X_m X_{m-2} \\
&= E \sum_m X_m X_{m-1} - \sum_m X_{m-1}^2 - \sum_m X_{m+1} X_{m-1} \\
&= \sum_m X_{m-1} (EX_m - X_{m-1} - X_{m+1}) \\
&= \lambda \sum_m X_m X_{m-1} C_m.
\end{aligned} \tag{2.3}$$

Subtracting the right-hand side from the left-hand side in (2.3), we get

$$\sum_m X_m X_{m-1} \sin(2m-1)\pi\theta = 0. \tag{2.4}$$

By (2.2) we infer that

$$\begin{aligned} E^2 &= 4+\lambda^2 - \sum_m (X_{m+1}-X_{m-1}+\lambda X_m \sin 2m\pi\theta)^2 \\ &\quad + 2\lambda \sum_m X_m X_{m-1}(C_m + C_{m-1}) + 2\lambda \sum_m X_{m+1}X_m \sin 2m\pi\theta \\ &\quad - 2\lambda \sum_m X_m X_{m-1} \sin 2m\pi\theta \\ &= 4+\lambda^2 - \sum_m (X_{m+1}-X_{m-1}+\lambda X_m \sin 2m\pi\theta)^2 \\ &\quad + 2\lambda \sum_m X_m X_{m-1}\big(C_m + C_{m-1} + \sin 2(m-1)\pi\theta - \sin 2m\pi\theta\big). \end{aligned} \tag{2.5}$$

But

$$\begin{aligned} &\cos 2m\pi\theta + \cos 2(m-1)\pi\theta + \sin 2(m-1)\pi\theta - \sin 2m\pi\theta \\ &\quad = \cos 2m\pi\theta + \cos 2m\pi\theta \cos 2\pi\theta + \sin 2m\pi\theta \sin 2\pi\theta \\ &\qquad + \sin 2m\pi\theta \cos 2\pi\theta - \cos 2m\pi\theta \sin 2\pi\theta - \sin 2m\pi\theta \\ &\quad = \cos 2m\pi\theta(1+\cos 2\pi\theta - \sin 2\pi\theta) - \sin 2m\pi\theta(1-\cos 2\pi\theta - \sin 2\pi\theta) \\ &\quad = 2\cos 2m\pi\theta \cos \pi\theta(\cos\pi\theta - \sin\pi\theta) - 2\sin 2m\pi\theta \sin\pi\theta(\sin\pi\theta - \cos\pi\theta) \\ &\quad = 2(\cos\pi\theta - \sin\pi\theta)(\cos\pi\theta \cos 2m\pi\theta + \sin\pi\theta \sin 2m\pi\theta) \\ &\quad = 2(\cos\pi\theta - \sin\pi\theta)\cos(2m-1)\pi\theta, \end{aligned}$$

which gives in conjunction with (2.5)

$$\begin{aligned} E^2 &= 4+\lambda^2 - \sum_m (X_{m+1}-X_{m-1}+\lambda X_m \sin 2m\pi\theta)^2 \\ &\quad + 4\lambda(\cos\pi\theta - \sin\pi\theta) \sum_m X_m X_{m-1} \cos(2m-1)\pi\theta. \end{aligned} \tag{2.6}$$

Using also

$$C_m + C_{m-1} = \cos 2m\pi\theta + \cos 2(m-1)\pi\theta = 2\cos\pi\theta \cos(2m-1)\pi\theta,$$

we derive from (2.6) and (2.3)

$$\begin{aligned} E^2 &= 4+\lambda^2 - \sum_m (X_{m+1}-X_{m-1}+\lambda X_m \sin 2m\pi\theta)^2 \\ &\quad + 4\lambda(1-\tan\pi\theta) \sum_m X_m X_{m-1} C_m \qquad 0 \le \theta < 1/2. \end{aligned} \tag{2.7}$$

From (2.6) it follows that

$$E^2 \le 4+\lambda^2 + 4\lambda(\cos\pi\theta - \sin\pi\theta) \sum_m X_m X_{m-1} \cos(2m-1)\pi\theta. \tag{2.8}$$

Using the identity

$$2(ax+by) = (a+b)(x+y) + (a-b)(x-y)$$

for $a = C_{m+1}$, $b = C_{m-1}$, $x = X_{m+1}$, $y = X_{m-1}$, we may write

$$\begin{aligned} 2\lambda \sum_m & X_m X_{m-1}(C_m + C_{m-1}) \\ &= \lambda \sum_m X_m X_{m-1}(C_m + C_{m-1}) + \lambda \sum_m X_{m+1} X_m (C_{m+1} + C_m) \\ &= \lambda \sum_m X_m (X_{m+1} + X_{m-1}) C_m \\ &\qquad + \frac{\lambda}{2} \sum_m X_m (X_{m+1} + X_{m-1})(C_{m+1} + C_{m-1}) \\ &\qquad + \frac{\lambda}{2} \sum_m X_m (X_{m+1} - X_{m-1})(C_{m+1} - C_{m-1}) \\ &= \frac{\lambda}{2} \sum_m X_m (X_{m+1} + X_{m-1})(C_{m+1} + 2C_m + C_{m-1}) \\ &\qquad + \frac{\lambda}{2} \sum_m X_m (X_{m+1} - X_{m-1})(C_{m+1} - C_{m-1}). \end{aligned} \tag{2.9}$$

Since

$$C_{m+1} - C_{m-1} = \cos 2(m+1)\pi\theta - \cos 2(m-1)\pi\theta = -2 \sin 2\pi\theta \sin 2m\pi\theta,$$

we collect from (2.2) and (2.9)

$$\begin{aligned} E^2 = 4 + \lambda^2 &+ \frac{\lambda}{2} \sum_m X_m (X_{m+1} + X_{m-1})(C_{m+1} + 2C_m + C_{m-1}) \\ &- \lambda^2 \sum_m X_m^2 \sin^2 2m\pi\theta - \sum_m (X_{m+1} - X_{m-1})^2 \\ &- \lambda \sin 2\pi\theta \sum_m X_m (X_{m+1} - X_{m-1}) \sin 2m\pi\theta. \end{aligned} \tag{2.10}$$

Using

$$\begin{aligned} \sum_m & X_m (X_{m+1} + X_{m-1})(C_{m+1} + 2C_m + C_{m-1}) \\ &= \sum_m \Big(X_{m-1} X_m (C_m + 2C_{m-1} + C_{m-2}) + X_{m-1} X_m (C_{m+1} + 2C_m + C_{m-1}) \Big) \\ &= \sum_m X_m X_{m-1} (C_{m+1} + 3C_m + 3C_{m-1} + C_{m-2}), \end{aligned}$$

$$\begin{aligned} C_{m+1} + 3C_m + 3C_{m-1} + C_{m-2} & \\ &= \Re(e^{2\pi i(m+1)\theta} + 3e^{2\pi i m\theta} + 3e^{2\pi i(m-1)\theta} + e^{2\pi i(m-2)\theta}) \\ &= \Re\big(e^{2\pi i(m-2)\theta}(1 + e^{2\pi i\theta})^3\big) = \Re\big(e^{(2m-1)\pi i\theta}(e^{-\pi i\theta} + e^{\pi i\theta})^3\big) \\ &= \Re\big(e^{(2m-1)\pi i\theta}(2\cos\pi\theta)^3\big) = 8\cos^3\pi\theta \cos(2m-1)\pi\theta \end{aligned}$$

and (2.10), we collect

$$\begin{aligned}
E^2 &= 4+\lambda^2+4\lambda\cos^3\pi\theta\sum_m X_mX_{m-1}\cos(2m-1)\pi\theta \\
&\quad -\lambda^2\sum_m X_m^2\sin^2 2m\pi\theta-\sum_m(X_{m+1}-X_{m-1})^2 \\
&\quad -\lambda\sin 2\pi\theta\sum_m X_m(X_{m+1}-X_{m-1})\sin 2m\pi\theta \\
&= 4+\lambda^2+4\lambda\cos^3\pi\theta\sum_m X_mX_{m-1}\cos(2m-1)\pi\theta \\
&\quad -\frac{\lambda^2}{4}(4-\sin^2 2\pi\theta)\sum_m X_m^2\sin^2 2m\pi\theta \\
&\quad -\sum_m\left(\frac{\lambda}{2}X_m\sin 2\pi\theta\sin 2m\pi\theta+X_{m+1}-X_{m-1}\right)^2 \\
&\le 4+\lambda^2+4\lambda\cos^3\pi\theta\sum_m X_mX_{m-1}\cos(2m-1)\pi\theta.
\end{aligned} \tag{2.11}$$

We can now prove

Theorem 2.1. (i) *For every $\theta\in[1/4,1/2]$, we have*

$$\|H_{\theta,\lambda}\|\le\sqrt{4+\lambda^2}.$$

(ii) *For every $\theta\in[0,1/4]$, we have*

$$\|H_{\theta,\lambda}\|^2\le 4+\lambda^2+4\lambda(\cos\pi\theta-\sin\pi\theta)\cos\pi\theta.$$

Proof. Since $H_{\theta,\lambda}$ and $H_{\theta,-\lambda}$ have the same spectrum, we may assume that $\lambda>0$. Using the continuity of the map $\theta\mapsto\|H_{\theta,\lambda}\|$, it suffices to assume in the first place that θ is rational. Taking now stock on (1.1), we may replace $\|H_{\theta,\lambda}\|$ by $\|h_{\theta,\lambda}\|$.

(i) Let $\theta\in[1/4,1/2)$. The sum $\sum_m X_mX_{m-1}\cos(2m-1)\pi\theta$ played a central role in the previous computations. We do not have any control over this sum. However, the point is that if $\sum_m X_mX_{m-1}\cos(2m-1)\pi\theta\le 0$, then (2.11) gives $E^2\le 4+\lambda^2$, and if $\sum_m X_mX_{m-1}\cos(2m-1)\pi\theta\ge 0$, then (2.8) gives $E^2\le 4+\lambda^2$.

(ii) This estimate follows from (2.8) and from the next lemma. □

Lemma 2.2. *For every $\theta\in[0,1/2]$, we have*

$$\begin{aligned}
\left|\sum_m X_mX_{m-1}\cos(2m-1)\pi\theta\right| &\le \frac{1}{2}\sqrt{2(1+|\cos 2\pi\theta|)} \\
&= \begin{cases}\cos\pi\theta & \text{if } 0\le\theta\le 1/4\\ \sin\pi\theta & \text{if } 1/4\le\theta\le 1/2.\end{cases}
\end{aligned}$$

Proof. With $D_m = \cos(2m-1)\pi\theta$, we gather

$$\begin{aligned}&D_m^2 + D_{m+1}^2 = \cos^2(2m-1)\pi\theta + \cos^2(2m+1)\pi\theta\\ &\quad = 1 + \big(\cos(4m-2)\pi\theta + \cos(4m+2)\pi\theta\big)/2 = 1 + \cos 4m\pi\theta \cos 2\pi\theta\\ &\le 1 + |\cos 2\pi\theta|.\end{aligned}$$

Combining this with Cauchy-Schwartz, we derive

$$\begin{aligned}2\sum_m |X_m X_{m-1} D_m| &= \sum_m \big(|X_m||X_{m-1}||D_m| + |X_{m+1}||X_m||D_{m+1}|\big)\\ &= \sum_m |X_m|\big(|X_{m-1}||D_m| + |X_{m+1}||D_{m+1}|\big)\\ &\le \sum_m |X_m|\sqrt{X_{m-1}^2 + X_{m+1}^2}\,\sqrt{D_m^2 + D_{m+1}^2}\\ &\le \sqrt{1+|\cos 2\pi\theta|}\,\sum_m |X_m|\,\sqrt{X_{m-1}^2 + X_{m+1}^2}\\ &\le \sqrt{1+|\cos 2\pi\theta|}\,\sqrt{\sum_m X_m^2}\,\sqrt{\sum_m (X_{m-1}^2 + X_{m+1}^2)}\\ &= \sqrt{2(1+|\cos 2\pi\theta|)}.\end{aligned}$$

□

3. Improved upper bound estimates in the range $\theta \in [1/4, 1/2]$

Since the spectra of $H_{\theta,\lambda}$ and $H_{\theta,-\lambda}$ coincide, we may assume that $\lambda > 0$. We also assume throughout this section that $\theta \in (1/4, 1/2)$.

Using the notation from the previous section, we set

$$\begin{aligned}S = S(\theta,\lambda,E) &= E\sum_m C_m X_m^2 + \frac{2}{\lambda}\sum_m X_{m+1}X_{m-1},\\ T = T(\theta,\lambda) &= \lambda\sum_m C_m^2 X_m^2 + \frac{2}{\lambda}\sum_m X_{m+1}X_{m-1}.\end{aligned}$$

It is worth to note first that

$$\begin{aligned}\sum_m X_m X_{m-1} = \sum_m X_{m+1}X_m &= \sum_m \big((E-\lambda C_m)X_m - X_{m-1}\big)X_m\\ &= E - \lambda\sum_m C_m X_m^2 - \sum_m X_m X_{m-1},\end{aligned}$$

thus

$$\sum_m X_m X_{m-1} = \sum_m X_{m+1}X_m = \frac{E}{2} - \frac{\lambda}{2}\sum_m C_m X_m^2, \tag{3.1}$$

and as a result

$$\begin{aligned}
2\lambda\sum_m X_m X_{m-1} C_m &= \lambda\sum_m (X_{m+1}X_m C_{m+1} + X_m X_{m-1} C_m) \\
\text{by (2.3)} \quad &= \lambda\sum_m X_{m+1}X_m C_m + \lambda\sum_m X_m X_{m-1} C_m \\
&= \lambda\sum_m C_m X_m (X_{m+1} + X_{m-1}) \\
\text{by (2.1)} \quad &= \sum_m (EX_m - X_{m+1} - X_{m-1})(X_{m+1} + X_{m-1}) \\
&= 2E\sum_m X_{m+1}X_m - \sum_m (X_{m+1} + X_{m-1})^2 \\
\text{by (3.1)} \quad &= 2E\left(\frac{E}{2} - \frac{\lambda}{2}\sum_m C_m X_m^2\right) - 2 - 2\sum_m X_{m+1}X_{m-1} \\
&= E^2 - \lambda E\sum_m C_m X_m^2 - 2\sum_m X_{m+1}X_{m-1} - 2 \\
&= E^2 - \lambda S - 2.
\end{aligned} \tag{3.2}$$

Therefore, (2.7) implies

$$E^2 \leq 4 + \lambda^2 + 2(1 - \tan\pi\theta)(E^2 - \lambda S - 2),$$

and so,

$$E^2 \leq \frac{\lambda^2 + 2\lambda(\tan\pi\theta - 1)S + 4\tan\pi\theta}{2\tan\pi\theta - 1},$$

which gives in turn

$$E^2 - 4 - \lambda^2 \leq \frac{2(\tan\pi\theta - 1)(\lambda S - \lambda^2 - 2)}{2\tan\pi\theta - 1}. \tag{3.3}$$

Lemma 3.1. *Suppose that* $(Y_m)_{m\in\mathbb{Z}_q}$ *is a unit vector in* $\ell^2(\mathbb{Z}_q)$. *Then*

$$2\sum_m C_m^2 Y_m^2 + \sum_m Y_{m+1}Y_{m-1} \leq 1 + \sqrt{2(1 + \cos^2 4\pi\theta)}.$$

Proof. If we write

$$2\sum_m C_m^2 Y_m^2 = 1 + \sum_m Y_m^2 \cos 4m\pi\theta,$$

then the desired result will follow at once as we have shown that

$$\sup_m \frac{Y_{m+1}^2 \cos 4(m+1)\pi\theta + Y_{m-1}^2 \cos 4(m-1)\theta + 2Y_{m+1}Y_{m-1}}{Y_{m+1}^2 + Y_{m-1}^2} \tag{3.4}$$

is less or equal than $\sqrt{2(1+\cos^2 4\pi\theta)}$. Taking $A = Y_{m+1}/\sqrt{Y_{m+1}^2 + Y_{m-1}^2}$ and $B = Y_{m-1}/\sqrt{Y_{m+1}^2 + Y_{m-1}^2}$, the expression in (3.4) is no greater than

$$\begin{aligned}
\sup_{\substack{m\in\mathbb{Z}\\ A^2+B^2=1}} & \Big((A^2+B^2)\cos 4\pi\theta \cos 4m\pi\theta + (B^2-A^2)\sin 4\pi\theta \sin 4m\pi\theta + 2AB \Big) \\
& = \sup_{A^2+B^2=1} \Big(\sqrt{(A^2+B^2)^2\cos^2 4\pi\theta + (B^2-A^2)^2 \sin^2 4\pi\theta} + 2AB \Big) \\
& \quad = \sup_{A^2+B^2=1} \Big(\sqrt{1-4A^2B^2\sin^2 4\pi\theta} + 2AB \Big) .
\end{aligned}$$

The statement follows now from

$$\begin{aligned}
\big(\sqrt{1-4A^2B^2\sin^2 4\pi\theta} + 2AB \big)^2 & \leq 2(1-4A^2B^2\sin^2 4\pi\theta + 4A^2B^2) \\
& = 2(1+4A^2B^2\cos^2 4\pi\theta) \leq 2(1+\cos^2 4\pi\theta).
\end{aligned}$$

□

Corollary 3.2. *If* $(Y_m)_{m\in\mathbb{Z}_q}$ *is a unit vector in* $\ell^2(\mathbb{Z}_q)$ *and* $0 \leq \lambda \leq 2$, *then*

$$\lambda \sum_m C_m^2 Y_m^2 + \frac{2}{\lambda} \sum_m Y_{m+1}Y_{m-1} \leq \frac{2}{\lambda} + \lambda \sqrt{\frac{1+\cos^2 4\pi\theta}{2}} .$$

Proof. Using lemma 3.1 and Cauchy-Schwarz, the expression in the left-hand side becomes

$$\begin{aligned}
\frac{\lambda}{2} & \Big(2\sum_m C_m^2 Y_m^2 + \sum_m Y_{m+1}Y_{m-1} \Big) + \Big(\frac{2}{\lambda} - \frac{\lambda}{2} \Big) \sum_m Y_{m+1}Y_{m-1} \\
& \leq \frac{2}{\lambda} - \frac{\lambda}{2} + \frac{\lambda}{2} \big(1 + \sqrt{2(1+\cos^2 4\pi\theta)} \big) = \frac{2}{\lambda} + \lambda \sqrt{\frac{1+\cos^2 4\pi\theta}{2}} .
\end{aligned}$$

□

Theorem 3.3. *For every* $\theta \in [1/4, 1/2]$ *and* $\lambda \in \mathbb{R}$, *we have*

$$\| H_{\theta,\lambda} \| \leq \sqrt{ 4 + \lambda^2 - \Big(1 - \frac{1}{\tan \pi\theta} \Big) \Big(1 - \sqrt{\frac{1+\cos^2 4\pi\theta}{2}} \Big) \min(4, \lambda^2) } .$$

Proof. We may assume that $\lambda > 0$. We may also assume as in the proof of Theorem 2.1 that θ is rational and then replace $\|H_{\theta,\lambda}\|$ by $\|h_{\theta,\lambda}\|$. Let ε be a constant to be chosen later.

We first consider the case $\lambda \leq 2$. Suppose that

$$2\lambda \sum_m X_m X_{m-1} C_m \leq \varepsilon. \tag{3.5}$$

Since

$$\begin{aligned}2\sum_m X_mX_{m-1}C_m &= \sum_m X_mX_{m-1}C_m + \sum_m X_{m+1}X_mC_m\\ &= \sum_m C_mX_m(X_{m-1}+X_{m+1}) = \sum_m (E-\lambda C_m)C_mX_m^2\\ &= E\sum_m C_mX_m^2 - \lambda\sum_m C_m^2X_m^2\\ &= S-T,\end{aligned}$$

we infer from (3.5) that

$$S \le \frac{\varepsilon}{\lambda} + T. \tag{3.6}$$

We combine (3.6) with (3.3) to collect

$$E^2-4-\lambda^2 \le \frac{2(\tan\pi\theta-1)}{2\tan\pi\theta-1}\,(\varepsilon+\lambda T-\lambda^2-2). \tag{3.7}$$

If

$$2\lambda\sum_m X_mX_{m-1}C_m \ge \varepsilon,$$

then we infer from (2.7) and from $1-\tan\pi\theta \le 0$ that

$$E^2-4-\lambda^2 \le 4\lambda(1-\tan\pi\theta)\sum_m X_mX_{m-1}C_m \le -2(\tan\pi\theta-1)\varepsilon. \tag{3.8}$$

In summary, we combine (3.7) with (3.8) to get

$$E^2-4-\lambda^2 \le 2(\tan\pi\theta-1)\max\Bigg(-\varepsilon, \frac{\varepsilon+\lambda T-\lambda^2-2}{2\tan\pi\theta-1}\Bigg)$$

for any $\varepsilon\in\mathbb{R}$. The inequality

$$\|H_{\theta,\lambda}\| \le \sqrt{4+\lambda^2-\left(1-\frac{1}{\tan\pi\theta}\right)\left(1-\sqrt{\frac{1+\cos^2 4\pi\theta}{2}}\right)\lambda^2} \tag{3.9}$$

follows by chosing

$$\varepsilon = \frac{\lambda^2+2-\lambda T}{2\tan\pi\theta}$$

and employing Corollary 3.2.

When $\lambda>2$, the desired inequality is obtained combining (3.9) with the Andre-Aubry duality

$$\|H_{\theta,\lambda}\| = \frac{\lambda}{2}\,\|H_{\theta,\frac{4}{\lambda}}\|.$$

□

4. SOME LOWER BOUNDS FOR THE NORM OF A HARPER OPERATOR

We shall consider sequences $(x_n)_n$ and $(y_n)_n$ in ℓ^2, of the form

$$x_n = x_n(\theta) = r^{|n|}$$

and

$$y_n = y_n(\theta) = \begin{cases} Ar^{|k|}, & \text{if } n = 2k \\ Br^{|k|}, & \text{if } n = 2k+1 \end{cases},$$

with $A^2 + B^2 = 1$, and with $0 < r = r(\theta) < 1$ to be chosen later. We have

$$\sum_n x_n^2 = \sum_n y_n^2 = \sum_n r^{2|n|} = \frac{1+r^2}{1-r^2}\,,$$

where we set $\sum\limits_n = \sum\limits_{n\in\mathbb{Z}}$.

Using the relations

$$\sum_k r^{|k|} \cos(ak+b) = \frac{(1-r^2)\cos b}{1-2r\cos a + r^2}\,, \tag{4.1}$$

$$\sum_k r^{|k|+|k-1|} \cos(ak+b) = \frac{2r(1-r^2)\cos\frac{a}{2}\cos\left(b+\frac{a}{2}\right)}{1-2r^2\cos a + r^4}\,, \tag{4.2}$$

and

$$\sum_k r^{|k+1|+|k-1|} = r^2 - 1 + \sum_k r^{2|k|},$$

we find

$$\begin{aligned} \|H_\theta\|^2 &\geq \frac{1-r^2}{1+r^2} \sum_n (2x_n \cos 2n\pi\theta + x_{n+1} + x_{n-1})^2 \\ &= \frac{1-r^2}{1+r^2} \sum_n \left(2r^{|n|}\cos 2n\pi\theta + r^{|n+1|} + r^{|n-1|}\right)^2 \\ &= 6 - \frac{(1-r^2)^2}{1+r^2} + \frac{2(1-r^2)^2}{1-2r^2\cos 4\pi\theta + r^4} + 8\,\frac{2r}{1+r^2}\cdot\frac{(1-r^2)^2\cos^2\pi\theta}{1-2r^2\cos 2\pi\theta + r^4}\,. \end{aligned}$$

Taking $r = \tan(\alpha/2)$ with $0 < \alpha < \pi/2$, it follows that $\|H_\theta\|^2$ is greater or equal than

$$\sup_{0<\alpha<\pi/2} \left(6 - \frac{4\cos^2\alpha}{1+\cos\alpha} + 2\cos^2\alpha\left(\frac{1}{1-\sin^2\alpha\cos^2 2\pi\theta} + \frac{4\sin\alpha\cos^2\pi\theta}{1-\sin^2\alpha\cos^2\pi\theta}\right)\right).$$

Chosing α such that

$$\cos^2\alpha = \frac{\sin\pi\theta}{1+\sin\pi\theta}\,,$$

we arrive at

$$\|H_\theta\|^2 \geq f_1(\theta)^2 \qquad \theta \in [0, 1/2], \tag{4.3}$$

where $f_1(\theta)$ is defined in (1.10).

In particular, we infer from (4.3) and (1.10) that

$$\min_{1/4\le\theta\le 1/2}\|H_\theta\| \ge \min_{1/4<\theta<1/2} f_1(\theta) \approx \sqrt{6.59303} \approx 2.56769. \tag{4.4}$$

To get a better estimate in a neighborhood of 1/4 and 1/2, we use the sequence $(y_n)_n$ and relations (4.1) and (4.2), to collect

$$\begin{aligned}
\sum_n (2y_n\cos 2n\pi\theta + y_{n+1} + y_{n-1})^2 &= \sum_k \Big(2Ar^{|k|}\cos 4k\pi\theta + B(r^{|k|} + r^{|k-1})\Big)^2 \\
&\quad + \sum_k \Big(2Br^{|k|}\cos(4k+2)\pi\theta + A(r^{|k|} + r^{|k+1|})\Big)^2 \\
&= 4A^2\sum_k r^{2|k|}\cos^2 4k\pi\theta + 4B^2\sum_k \cos^2(4k+2)\pi\theta \\
&\quad + 2(A^2+B^2)\sum_k (r^{2|k|} + r^{|k|+|k-1|}) \\
&\quad + 4AB\sum_k r^{2|k|}\big(\cos 4k\pi\theta + \cos(4k+2)\pi\theta\big) \\
&\quad + 4AB\sum_k r^{|k|+|k-1|}\big(\cos 4k\pi\theta + \cos(4k-2)\pi\theta\big).
\end{aligned}$$

This further equals

$$\begin{aligned}
&4\sum_k r^{2|k|} + 2A^2\sum_k r^{2|k|}\cos 8k\pi\theta + 2B^2\sum_k r^{2|k|}\cos(8k+4)\pi\theta \\
&\qquad + 2(A^2+B^2)\sum_k r^{|k|+|k-1|} + 8AB\cos\pi\theta\sum_k r^{2|k|}\cos(4k+1)\pi\theta \\
&\qquad + 8AB\cos\pi\theta\sum_k r^{|k|+|k-1|}\cos(4k-1)\pi\theta \\
&= 4\,\frac{1+r^2}{1-r^2} + 2A^2\,\frac{1-r^4}{1-2r^2\cos 8\pi\theta + r^4} + 2B^2\,\frac{(1-r^4)\cos 4\pi\theta}{1-2r^2\cos 8\pi\theta + r^4} \\
&= \frac{4r}{1-r^2} + \frac{8AB\cos^2\pi\theta(1-r^4)}{1-2r^2\cos 4\pi\theta} \\
&\qquad + 8AB\cos^2\pi\theta\cos 2\pi\theta\cdot\frac{2r}{1-r^2}\cdot\frac{1-r^4}{1-2r^2\cos 4\pi\theta + r^4}\,.
\end{aligned}$$

Hence

$$\begin{aligned}
\|H_\theta\|^2 &\ge 4 + \frac{4r}{1+r^2} + \frac{2(A^2+B^2\cos 4\pi\theta)(1-r^2)^2}{1-2r^2\cos 8\pi\theta + r^4} \\
&\qquad + 8AB\cos^2\pi\theta\left(1 + \frac{2r}{1+r^2}\,\cos 2\pi\theta\right)\frac{(1-r^2)^2}{1-2r^2\cos 4\pi\theta + r^4}\,.
\end{aligned}$$

Taking $r = \tan(\alpha/2)$ with $0 < \alpha < \pi/2$, we find

$$\begin{aligned} \|H_\theta\|^2 \geq 4 + 2\sin\alpha + \frac{2(A^2 + B^2 \cos 4\pi\theta)\cos^2\alpha}{1 - \sin^2\alpha\cos^2 4\pi\theta} \\ + 8AB\cos^2\pi\theta(1 + \sin\alpha\cos 2\pi\theta)\,\frac{\cos^2\alpha}{1 - \sin^2\alpha\cos^2 2\pi\theta}\,. \end{aligned} \tag{4.5}$$

Taking also

$$\sin^2\alpha = \frac{1 - |\sin 4\pi\theta|}{\cos^2 4\pi\theta} = \frac{1}{1 + |\sin 4\pi\theta|}\,, \qquad \cos^2\alpha = \frac{|\sin 4\pi\theta|}{1 + |\sin 4\pi\theta|}\,,$$

we get

$$\frac{\cos^2\alpha}{1 - \sin^2\alpha\cos^2 4\pi\theta} = \frac{1}{1 + |\sin 4\pi\theta|}$$

and

$$\frac{\cos^2\alpha}{1 - \sin^2\alpha\cos^2 2\pi\theta} = \frac{2|\cos 2\pi\theta|}{2|\cos 2\pi\theta| + |\sin 2\pi\theta|} = \frac{2}{2 + |\tan 2\pi\theta|}\,.$$

Hence (4.5) gives

$$\begin{aligned} \|H_\theta\|^2 \geq 4 + \frac{2}{\sqrt{1 + |\sin 4\pi\theta|}} + \frac{2(A^2 + B^2\cos 4\pi\theta)}{1 + |\sin 4\pi\theta|} \\ + \frac{16AB\cos^2\pi\theta}{2 + |\tan 2\pi\theta|}\left(1 + \frac{\cos 2\pi\theta}{\sqrt{1 + |\sin 4\pi\theta|}}\right). \end{aligned} \tag{4.6}$$

Using also the equality

$$\max_{A^2+B^2=1} 2(\alpha_0 A^2 + \beta_0 B^2 + \gamma_0 AB) = \alpha_0 + \beta_0 + \sqrt{(\alpha_0 - \beta_0)^2 + \gamma_0^2}, \tag{4.7}$$

we infer from (4.6), with

$$\begin{aligned} \alpha_0 = \frac{1}{1 + |\sin 4\pi\theta|}\,, \qquad \beta_0 = \frac{\cos 4\pi\theta}{1 + |\sin 4\pi\theta|} \\ \gamma_0 = \frac{8\cos^2\pi\theta}{2 + |\tan 2\pi\theta|}\left(1 + \frac{\cos 2\pi\theta}{\sqrt{1 + |\sin 4\pi\theta|}}\right), \end{aligned}$$

that

$$\|H_\theta\|^2 \geq f_2(\theta)^2 \qquad 0 \leq \theta \leq 1/2, \tag{4.8}$$

with $f_2(\theta)$ given by (1.11).

If we take

$$z_n = \begin{cases} \sqrt{1/10}\,\sin\alpha & \text{if } n = \pm 2 \\ \sqrt{2/5}\,\sin\alpha & \text{if } n = \pm 1 \\ \cos\alpha & \text{if } n = 0 \\ 0 & \text{if } n \neq 0, \pm 1, \pm 2, \end{cases}$$

for some $\alpha \in \mathbb{R}$, then $\sum_n z_n^2 = 1$, and so we get

$$\begin{aligned}\|H_\theta\|^2 &\geq 2\bigg(\frac{\sin\alpha}{\sqrt{10}}\bigg)^2 + 2\bigg(\frac{2\cos 4\pi\theta \sin\alpha}{\sqrt{10}} + \sqrt{\frac{2}{5}}\sin\alpha\bigg)^2 \\ &= 2\bigg(2\cos 2\pi\theta\sqrt{\frac{2}{5}}\sin\alpha + \cos\alpha + \frac{\sin\alpha}{\sqrt{10}}\bigg)^2 + \bigg(2\cos\alpha + 2\sqrt{\frac{2}{5}}\sin\alpha\bigg)^2.\end{aligned}$$

Since the right-hand side of the inequality above is equal to

$$\begin{aligned}&\frac{\sin^2\alpha}{5} + \frac{4}{5}\sin^2\alpha(1+\cos 4\pi\theta)^2 + 2\bigg(\cos\alpha + \frac{\sin\alpha(1+4\cos 2\pi\theta)}{\sqrt{10}}\bigg)^2 \\ &\qquad + 4\bigg(\cos\alpha + \sqrt{\frac{2}{5}}\sin\alpha\bigg)^2 \\ &\quad = 6\cos^2\alpha + \frac{\sin^2\alpha(9 + 16\cos^4 2\pi\theta + (1+4\cos 2\pi\theta)^2)}{5} \\ &\qquad + \sin 2\alpha\bigg(\frac{2(1+4\cos 2\pi\theta)}{\sqrt{10}} + 4\sqrt{\frac{2}{5}}\bigg) \\ &\quad = 6\cos^2\alpha + 2\sin^2\alpha\bigg(1 + \frac{4}{5}(\cos 2\pi\theta + 2\cos^2 2\pi\theta + 2\cos^4 2\pi\theta)\bigg) \\ &\qquad + \sin 2\alpha\bigg(\sqrt{10} + \frac{8\cos 2\pi\theta}{\sqrt{10}}\bigg),\end{aligned}$$

we deduce from (4.7) that

$$\|H_\theta\|^2 \geq f_3(\theta)^2, \tag{4.9}$$

where $f_3(\theta)$ is as in (1.12) .

References

1. J. AVRON, P. H. M. van MOUCHE and B. SIMON, On the measure of the spectrum for the almost Mathieu operator, *Comm. Math. Phys.* 132 (1990), 103–118.
2. C. BÉGUIN, A. VALETTE and A. ZUK, On the spectrum of a random walk on the discrete Heisenberg group and the norm of Harper's operator, *J. Geom. Phys.* 21 (1997), 337–356.
3. J. BELLISSARD, K-theory of C^*-algebras in solid state physics, *Statistical mechanics and field theory: mathematical aspects* (Groningen, 1985), Lecture Notes in Phys., 257, Springer, Berlin-New York, 1986, pp 99–156.
4. J. BELLISSARD, Lipshitz continuity of gap boundaries for Hofstadter-like spectra, *Comm. Math. Phys.* 160 (1994), 599-613.
5. J. BELLISSARD and B. SIMON, Cantor spectrum for the almost Mathieu equation, *J. Funct. Anal.* 48 (1982), 408–419.
6. F. P. BOCA. *Rotation C^*-algebras and almost Mathieu operators.* The Theta Foundation. Bucharest 2001.
7. J. BOURGAIN and M. GOLDSTEIN, On nonperturbative localization with quasi-periodic potential, *Ann. of Math.* 152 (2000), 835–879.
8. M. D. CHOI, G. A. ELLIOTT and N. YUI, Gauss polynomials and the rotation algebras, *Invent. Math.* 99 (1990), 225–246.

9. U. HAAGERUP and M. RØRDAM, Perturbations of the rotation C^*-algebras and of the Heisenberg commutation relations, *Duke Math. J.* 77 (1995), 627–656.
10. B. HELFFER and J. SJÖSTRAND, Analyse semi-classique pour l'équation de Harper.II. Comportement semi-classique près d'un rationnel, *Mém. Soc. Math. France* No.40 (1990).
    Semi-classical analysis for Harper's equation.III. Cantor structure of the spectrum, *Mém. Soc. Math. France (N.S.)* No.39 (1989).
11. D. R. HOFSTADTER, Energy levels and wave functions of Bloch electrons in a rational or irrational magnetic field, *Phys. Rev. B* 14 (1976), 2239–2249.
12. S. Ya. JITOMIRSKAYA, Metal-insulator transition for the almost Mathieu operator, *Ann. of Math.* 150 (1999), 1159–1175.
13. E. KIRCHBERG and N. C. PHILLIPS, Embedding of continuous fields of C^*-algebras in the Cuntz algebra $\mathcal{O}_2$, *J. Reine Angew. Mathematik* 525 (2000), 55–94.
14. Y. LAST, Zero measure spectrum for the almost Mathieu operator, *Comm. Math. Phys.* 164 (1994), 421–432.
15. R. SZWARC, Norm estimates of discrete Schrödinger operators, *Coll. Math.* 76 (1998), 153–160.

Department of Mathematics, University of Illinois, Urbana, IL 61801

Institute of Mathematics of the Romanian Academy, P.O.Box 1-764, Bucharest 70700, Romania

fboca@math.uiuc.edu zaharesc@math.uiuc.edu